

Microscopic Derivation of Ginzburg–Landau Theory and the BCS Critical Temperature Shift in a Weak Homogeneous Magnetic Field

Andreas Deuchert

Institute of Mathematics, University of Zurich (UZH)


Workshop II (Semiclassical Analysis, QFT, nonlinear PDE) @ Politecnico di Milano
May 26, 2022

Joint work with Christian Hainzl and Marcel Schaub

Funding from the European Union's Horizon 2020 research and innovation programme (Marie Skłodowska-Curie fellowship) and from the SNSF (Ambizione grant) is gratefully acknowledged.


Superfluidity

Vortex lattice of Fermi gas across the **BEC - BCS crossover**.




M. W. Zwierlein, J. R. Abo-Shaeer, A. Schirotzek, C. H. Schunck, W. Ketterle, *Vortices and superfluidity in a strongly interacting Fermi gas*, *Nature* **435**, 1047–1051 (2005)


Setup: Particles in a constant magnetic field



Setup: Particles in a constant magnetic field



Setup: Particles in a constant magnetic field



We consider **attractively** interacting fermionic particles in a weak external magnetic field $|\mathbf{B}| \ll 1$.

Fermionic quasi-free states

- States on the fermionic Fock space obeying the Wick theorem are called **quasi-free**.
- Such states are uniquely determined by their **generalized one-particle density matrix** $\Gamma \in \mathcal{L}(L^2(\mathbb{R}^3) \oplus L^2(\mathbb{R}^3))$, $0 \leq \Gamma \leq 1$, which is of the form

$$\Gamma = \begin{pmatrix} \gamma & \alpha \\ \bar{\alpha} & 1 - \bar{\gamma} \end{pmatrix}.$$

This, in particular, implies $0 \leq \gamma \leq 1$ and $\alpha\alpha^* \leq \gamma(1 - \gamma)$. We assume that $\alpha(x, y) = \alpha(y, x) \Rightarrow$ **s-wave Cooper pairs** (spin-singulett).

- Our states are **gauge-periodic** w.r.t. $\Lambda_B = \sqrt{2\pi B^{-1}}\mathbb{Z}^3$, that is, for $\lambda \in \Lambda_B$ we have

$$\gamma(x + \lambda, y + \lambda) = e^{-i\frac{\mathbf{B}}{2} \cdot (\lambda \wedge (x-y))} \gamma(x, y),$$

$$\alpha(x + \lambda, y + \lambda) = e^{-i\frac{\mathbf{B}}{2} \cdot (\lambda \wedge (x+y))} \alpha(x, y).$$

The BCS free energy functional

For gauge-periodic BCS states we define the **BCS free energy functional** by

$$\mathcal{F}_{\mathbf{B}, T}^{\text{BCS}}(\Gamma) = \text{Tr}[\left((-i\nabla + \mathbf{A})^2 - \mu\right)\gamma] - TS(\Gamma) - \frac{1}{|Q_B|} \int_{Q_B \times \mathbb{R}^3} V(x-y) |\alpha(x, y)|^2 dx dy,$$

where

- $\text{Tr}[A] = \frac{1}{|Q_B|} \text{Tr}_{L^2(Q_B)}[\mathbb{1}(x \in Q_B)A\mathbb{1}(x \in Q_B)]$, that is, we consider **energy per unit volume**, (Q_B is the **unit cell** of the lattice Λ_B)
- $\mathbf{A}(x) = \frac{1}{2}\mathbf{B} \wedge x$ (**Vector potential**),
- $\mu \in \mathbb{R}$, $T \geq 0$ (**chemical potential, temperature**),
- $S(\Gamma) = -\text{Tr}[\Gamma \ln(\Gamma)]$ (**Von Neumann entropy**),

Normal state, free energy and superconductivity

- The unique minimizer for $V = 0$ is called **normal state** and reads

$$\Gamma_0 = \begin{pmatrix} \gamma_0 & 0 \\ 0 & 1 - \bar{\gamma}_0 \end{pmatrix} \quad \text{with} \quad \gamma_0 = \frac{1}{e^{((-i\nabla + \mathbf{A}(x))^2 - \mu)/T} + 1}.$$

- The **BCS free energy** is defined by

$$F^{\text{BCS}}(B, T) = \inf_{\Gamma} \{ \mathcal{F}_{\mathbf{B}, T}^{\text{BCS}}(\Gamma) - \mathcal{F}_{\mathbf{B}, T}^{\text{BCS}}(\Gamma_0) \}.$$

- The system is said to be **superconducting** at temperature T and magnetic field \mathbf{B} if

$$F^{\text{BCS}}(B, T) < 0.$$

Effective models I: Translation-invariant BCS functional

If $\mathbf{B} = 0$ we describe the system by **translation-invariant states**, that is,

$$\gamma(x, y) = \gamma(x - y) \quad \text{and} \quad \alpha(x, y) = \alpha(x - y).$$

In this case it has been shown that there exists a **critical temperature** T_c such that the system is superconducting if $T < T_c$ and in its normal state if $T \geq T_c$.

Moreover, T_c can be defined as the **unique value of** T such that the operator

$$K_T - V(x) = \frac{-\Delta - \mu}{\tanh\left(\frac{-\Delta - \mu}{2T}\right)} - V(x)$$

has 0 as its lowest eigenvalue. The **corresponding eigenfunction** will be denoted by α_* .

Effective models II: The Ginzburg–Landau functional

We call a function $\Psi : \mathbb{R}^3 \rightarrow \mathbb{C}$ **gauge-periodic** if for all $\lambda \in \Lambda_B$ we have

$$\Psi(X + \lambda) = e^{-i\mathbf{B} \cdot (\lambda \wedge X)} \Psi(X).$$

For such functions and parameters $\Lambda_1, \Lambda_2, \Lambda_3 > 0$, $D \in \mathbb{R}$ the **Ginzburg–Landau functional** is defined by

$$\mathcal{E}_{\mathbf{B}, D}^{\text{GL}}(\Psi) = \frac{1}{B^2 |Q_B|} \int_{Q_B} (\Lambda_1 |(-i\nabla + 2\mathbf{A})\Psi(X)|^2 - DB\Lambda_2 |\Psi(X)|^2 + \Lambda_3 |\Psi(X)|^4) dX.$$

Its minimal value is the **Ginzburg–Landau energy**

$$E^{\text{GL}}(D) = \inf_{\Psi} \mathcal{E}_{\mathbf{B}, D}^{\text{GL}}(\Psi),$$

which, by scaling, does not depend on B .

Translation-invariant BCS functional

- Hainzl, Hamza, Seiringer, Solovej, *The BCS functional for general pair interactions*, CMP **281**, 349 (2008)
- Hainzl, Seiringer, *The Bardeen–Cooper–Schrieffer functional of superconductivity and its mathematical properties*, J. Math. Phys. **57**, 021101 (2016) (review)

BCS functional with external fields, zero magnetic flux through unit cell

- Frank, Hainzl, Seiringer, Solovej, *Microscopic derivation of Ginzburg–Landau theory*, J. Amer. Math. Soc. **25**, 667 (2012)
- Frank, Hainzl, Seiringer, Solovej, *The external field dependence of the BCS critical temperature*, CMP **342**, 189 (2016)

Linearized BCS theory, constant magnetic field

- Frank, Hainzl, Langmann, *The BCS critical temperature in a weak homogeneous magnetic field*, J. Spectr. Theory **9**, 1005 (2019)

Main result part I: Free energy asymptotics

Assumptions:

- V radial function with $(1 + |\cdot|^2)V \in L^\infty(\mathbb{R}^3)$,
- V such that $T_c > 0$ in translation-invariant case,
- Zero eigenvalue of $K_{T_c} - V(x)$ is simple.

Theorem (D., Hainzl, Schaub)

There are constants $C > 0$ and $B_0 > 0$ such that for all $0 < B \leq B_0$, we have

$$F_{\mathbf{B}, T_c(1-DB)}^{\text{BCS}} = B^2 (E^{\text{GL}}(D) + \mathcal{R})$$

with

$$-CB^{\frac{1}{12}} \leq \mathcal{R} \leq CB.$$

Main result part II: Approximate minimizers

Theorem (continued)

For any approximate minimizer Γ of $\mathcal{F}_{\mathbf{B},T}^{\text{BCS}}$ at $T = T_c(1 - DB)$ in the sense that

$$\mathcal{F}_{\mathbf{B},T}^{\text{BCS}}(\Gamma) - \mathcal{F}_{\mathbf{B},T}^{\text{BCS}}(\Gamma_0) \leq B^2 (E^{\text{GL}}(D) + \rho)$$

holds for some $\rho \geq 0$, we have the decomposition

$$\alpha(X, r) = \alpha_*(r)\Psi(X) + \sigma(X, r)$$


for the **Cooper pair wave function** $\alpha = \Gamma_{12}$. Here, σ satisfies

$$\frac{1}{|Q_B|} \int_{Q_B \times \mathbb{R}^3} |\sigma(X, r)|^2 d(X, r) \leq CB^{11/6},$$

α_* is the zero energy eigenfunction of $K_{T_c} - V(x)$, and Ψ obeys

$$\mathcal{E}_{\mathbf{B},D}^{\text{GL}}(\Psi) \leq E^{\text{GL}}(D) + \rho + CB^{1/12}.$$

Separation of scales



- Ψ approximate minimizer of $\mathcal{E}_{\mathbf{B},D}^{\text{GL}}$
- $(K_{T_c} - V(x))\alpha_* = 0$

Main result part III: Critical temperature shift

Assumptions: Same as for previous theorem.

Definition: $D_c = \Lambda_1 \inf \sigma_{L^2_{\text{mag}}(Q_1)}((-i\nabla + e_3 \wedge X)^2)$.

Theorem (D., Hainzl, Schaub)

There are constants $C > 0$ and $B_0 > 0$ such that for all $0 < B \leq B_0$ the following holds:

- Let $0 < T_0 < T_c$. If the temperature satisfies

$$T_0 \leq T \leq T_c(1 - B(D_c + CB^{1/2})),$$


then we have $F^{\text{BCS}}(B, T) < 0$.

- If the temperature satisfies


$$T \geq T_c(1 - B(D_c - CB^{1/12})),$$

then we have $\mathcal{F}_{B,T}^{\text{BCS}}(\Gamma) - \mathcal{F}_{B,T}^{\text{BCS}}(\Gamma_0) > 0$ unless $\Gamma = \Gamma_0$.

Phase Diagram



Phase Diagram



Main novelty

The main novelty of our work are the following a-priori bounds for low energy states:

Theorem (Structure of low energy states)

For all $D_0, D_1 \geq 0$, there is a constant $B_0 > 0$ such that for all $0 < B_0 \leq B$ we have the following statement: If $T - T_c \geq -D_0 B$ and if Γ obeys

$$\mathcal{F}_{\mathbf{B}, T}^{\text{BCS}}(\Gamma) - \mathcal{F}_{\mathbf{B}, T}^{\text{BCS}}(\Gamma_0) \leq D_1 B^2,$$

then there exist $\Psi \in H_{\text{mag}}^1(Q_B)$ and $\xi \in H_{\text{mag}}^1(Q_B \times \mathbb{R}^3)$ such that

$$\alpha(X, r) = \Psi(X)\alpha_*(r) + \xi(X, r),$$

where

$$\sup_{0 < B \leq B_0} \|\Psi\|_{H_{\text{mag}}^1(Q_B)}^2 \leq C \quad \text{and} \quad \|\xi\|_{H_{\text{mag}}^1(Q_B \times \mathbb{R}^3)}^2 \leq CB^2 \left(\|\Psi\|_{H_{\text{mag}}^1(Q_B)}^2 + D_1 \right).$$